

Albino Kingbird: On June 10, 1967, as we drove along Rte. 82 between Stanfordville and Washington Hollow in Dutchess County, a completely white robin-size bird flashed to our left. After stopping and observing the bird closely (15 feet at one time), Otis Waterman and I concluded it was a Kingbird. The bird had a yellow bill and legs, and dark eyes but otherwise was pure white. It made no sound though a normal Eastern Kingbird in the area did call.

Though I made several attempts to locate the bird again, it was not until July 6 that I received a phone call from the owner of the house in the yard of which we had first observed it, asking what the all-white bird might be. From then until August 14 the albino was seen almost daily by my husband. A number of others also had an opportunity to observe it, feeding in the air over a small marshy area and perching in low branches of wild cherry trees, preening and sunning itself. The bird was photographed on July 22 by Herbert Saltford and was last seen on August 27.

At no time was there any indication of its having been a breeding bird though a single normal Kingbird was often observed close by.

Florence Germond, Shunpike, Clinton Corners, N.Y. 12514

Ed. Note: There are not many reports of albino flycatchers, though a cream-colored Phoebe was reported near Etna Oct. 13, 1952 (*Kingbird* 2(4): 90, Dec. 1952).

Odd Warbler at Caton Swamp, near Elmira: On June 24, 1967, Louise Fivie and I observed an unusual warbler on the road through Caton Swamp southwest of Elmira. As we were driving along I heard the song of a Black-throated Green Warbler (*Dendroica virens*) close to the road and stopped so that I could point it out to Louise. The warbler flew to a sparsely needled branch of a large hemlock that arched the road directly in front of us. I was surprised to see a completely yellow head and immediately consulted my Eastern Field Guide, with no satisfaction. After singing the typical Black-throated Green song for approximately five minutes, the bird flew away.

Since he was facing us, I had no view of his back, but the front view showed no green. However, there were the typical black throat and white underparts. The light was excellent for viewing with 7×50 binoculars.

Later, the only bird that I could find that closely resembled this one was the Hermit Warbler (*Dendroica occidentalis*) of the western coast. Perhaps the bird was a hybrid Hermit-Black-throated Green, or it may have been Xanthochroism (excess of yellow in plumage).

Wilifred I. Howard, 610 Coburn Street, Elmira, N.Y. 14904

Immature Black-headed Gull at Sandy Pond Outlet, Oswego County: On July 22, 1967 at about 10:30 a.m. at the Sandy Pond outlet at the eastern end of Lake Ontario, C. G. Spies and I noted, with two Bonaparte's Gulls perched on a log in the outlet channel, a similar gull with a red bill. This gull was as much larger than the Bonaparte's as these were than Common Terns also on the log. A black spot behind the eye blended into a smudgy gray hood, the gray color extending forward to the eye and up from it across the top of the crown and back to the nape, where it cut off sharply. The head was otherwise white, as were the underparts. The tail was white except for the terminal 1/2 inch (about 1/5 the length of the tail beyond the coverts) which was black. As the tail was spread in preening, the black appeared to consist of a racket-shaped mark on the tip of each feather. The wings were the same gray as the wings of the Bonaparte's while the birds were perched. The feet were dark pink and the bill glowing dark coral red, seeming blackish on tip and ridge. The above markings were observed at about 400 ft. with 30× scope while the bird was perched, viewed against the water, part of the time preening its wings and tail.