

Wheatear at Van Cortlandt Park, New York City: At about 12.30 P.M. on June 8, 1968 near the Parade Ground in Van Cortlandt Park, we spotted a bird slightly smaller than a Hermit Thrust on the ground approximately 30 feet in front of us. As it moved constantly about in an open area with sparse tufts of grass, we could see little of it except the gray head with a black mask running through the eyes straight to the bill, lined with white above. It vaguely suggested a shrike—but the bill was short and pointed. As we moved closer, the bird hopped up a small incline, occasionally flashing white in its tail. When it emerged onto an open roadway, we could see that the back was dark blue-gray, the wings quite dark, almost black, the breast washed with creamy buff, the legs dark. We checked the legs for a band, in case it might be an escapee; there was no band. Despite the nervous twitching and wagging of its tail, the bird not only stayed on the roadway for most of five minutes, but also allowed us to approach fairly close to it along the road. Although one of us had seen Wheatears (*Oenanthe oenanthe*) several times in this full plumage in Europe, it did not occur to us that this bird was a Wheatear until it finally flew up into some nearby shrubbery, clearly showing us the black and white “inverted T” pattern on the tail. Shortly afterwards it disappeared and we were unable to find it again.

Ogden Goellet, Jr., 140 East 92nd Street, New York, N.Y. 10028

Susan Koelle, 157 East 81st Street, New York, N.Y. 10028

ED. NOTE: This is the eighth appearance of this species in the Marine Region. Two other records were also in the spring—May 13, 1956 at Orient and June 3, 1941 at Moriches Inlet (John Bull, *Birds of the New York Area*). Of interest is another Wheatear reported in nearby Connecticut this May (see future issue of *Audubon Field Notes* for details).
T. Davis

Unusual Concentration of Singing Slate-colored Juncos near Owego: On July 17, 1968 Mildred White and I observed an interesting number of singing Slate-colored Juncos (*Junco hyemalis*) in the Frost Hollow area near Owego. On four trips to this area, we found eight singing birds in a distance of about 2 miles. At one stop we had two very disturbed birds and one with food in its bill. Elevations range from 1000' to 1250'. In addition, on the cross road, about a quarter mile to the south we saw another junco that was tail-fanning, but found no other near. Along about one and a half miles of the northern section of this cross road we found four more singing birds. Elevations there are 1250' to 1350'. Hemlock is the predominant tree along both roads. Juncos regularly nest at these higher elevations, but we have not previously noted such a concentration in this area.

Ruth Williams, P.O. Box 382, Owego, N.Y. 13827

Apparent Recognition of Danger by Evening Grosbeaks: On July 17, 1967, I caught in a 2-cell Potter trap, and banded, two adult male Evening Grosbeaks (*Hesperiphona vespertina*) at the window feeder of Dora Burnett in Old Forge, New York. They were in the trap perhaps 15 minutes before being liberated.

A week later I set the trap in the same position, hoping to catch the four young that were reported there the day before, or a third unbanded male in the area. (Females had been seen earlier, but were not coming to the feeder with the young.) The males flew to the feeder from the near-by tree in which they usually perched, but refused to land. When a young one flew toward the trap, a male intercepted it and drove it to the ground, refusing to let it come to the feeder. Both birds then fed on the ground.

This behavior was repeated on July 27th, neither of the two banded males landing on the feeder, although they fed freely beneath it, and no young being allowed to land. A third, unbanded male was trapped successfully. All of these birds had been in the habit of eating at the feeder, rather than on the ground, so it would seem that the trap was recognized now as a danger.